



Pancreatic cancer and autoimmune diseases: An association sustained by computational and epidemiological case-control approaches

Journal:	<i>International Journal of Cancer</i>
Manuscript ID	IJC-18-1029.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Gómez-Rubio, Paulina; Spanish National Cancer Research Centre (CNIO), Genetic and Molecular Epidemiology Group; (2) Centro de Investigación Biomédica en Red en Oncología (CIBERONC)</p> <p>Piñero, Janet ; Research Program on Biomedical Informatics (GRIB), Hospital del Mar Research Institute (IMIM), Universidad Pompeu Fabra (UPF)</p> <p>Molina-Montes, Esther; Spanish National Cancer Research Centre (CNIO), Genetic and Molecular Epidemiology Group; (2) Centro de Investigación Biomédica en Red en Oncología (CIBERONC)</p> <p>Gutiérrez-Sacristán, Alba ; Research Program on Biomedical Informatics (GRIB), Hospital del Mar Research Institute (IMIM), Universidad Pompeu Fabra (UPF)</p> <p>Márquez, Mirari; Spanish National Cancer Research Centre (CNIO), Genetic and Molecular Epidemiology Group</p> <p>Rava, Marta; Spanish National Cancer Research Centre (CNIO), Genetic and Molecular Epidemiology Group</p> <p>Michalski, Christoph W. ; Technical University of Munich; University of Heidelberg</p> <p>Farré, Antoni; (6) Hospital de la Santa Creu i Sant Pau</p> <p>Molero, Xavier; (2) Centro de Investigación Biomédica en Red en Enfermedades Hepáticas y Digestivas (CIBERHD); Hospital Universitari Vall d'Hebron, Exocrine Pancreas Research Unit and Vall d'Hebron Research Institute (VHIR); Universitat Autònoma de Barcelona, Campus de la UAB</p> <p>Löhr, Matthias ; Karolinska Institutet and University Hospital, Gastrocentrum</p> <p>PEREA, JOSÉ; 12 DE OCTUBRE UNIVERSITY HOSPITAL,</p> <p>Greenhalf, Bill; University of Liverpool, MOLECULAR AND CLINICAL CANCER MEDICINE</p> <p>O'Rorke, Michael; Queen's University Belfast, Epidemiology & Public Health</p> <p>Tardon, Adonina; Universidad de Oviedo, Facultad de Medicina</p> <p>Gress, Thomas; University Hospital Marburg, Dept of Gastroenterology, Endocrinology and Metabolism</p> <p>Barberá, Victor M.; Hospital General Universitario de Elche</p> <p>Crnogorac-Jurcevic, Tatjana; Barts and The London School of Medicine and Dentistry, Molecular Pathology Unit</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	Muñoz-Bellví, Luís; Hospital Universitario de Salamanca Domínguez-Muñoz, Enrique ; Hospital Clínico Universitario de Santiago de Compostela Balsells, Joaquim ; (2) Centro de Investigación Biomédica en Red en Enfermedades Hepáticas y Digestivas (CIBERHD); Hospital Universitari Vall d´Hebron, Exocrine Pancreas Research Unit and Vall d´Hebron Research Institute (VHIR); Universitat Autònoma de Barcelona, Campus de la UAB Costello, Eithne; University of Liverpool, Department of Surgery Yu , Jingru ; Karolinska Institutet and University Hospital, Gastrocentrum Iglesias, Mar; Hospital del Mar/Parc de Salut Mar Ilzarbe, Lucas ; Hospital del Mar/Parc de Salut Mar Kleeff, Jörg ; Technical University of Munich; Martin-Luther-Universität Halle Wittenberg Juristische und Wirtschaftswissenschaftliche Fakultät Kong, Bo; Technical University of Munich Mora, Josefina ; (6) Hospital de la Santa Creu i Sant Pau Murray, Liam; Queen's University, Centre for Clinical and Population Sciences O'Driscoll, Damian ; National Cancer Registry Ireland Poves, Ignasi ; Hospital del Mar/Parc de Salut Mar Lawlor, Rita T.; University Hospital Trust of Verona Ye, Weimin; Karolinska Institute, Department of Medical Epidemiology; Hidalgo, Manuel; Beth Israel Deaconess Medical Center, Hematology / Oncology, Pancreatic Cancer; Scarpa, Aldo; Università di Verona, Dept. Pathology and Diagnostics; Sharp, Linda; National Cancer Registry Ireland; Institute of Health and Society, Newcastle University Carrato, Alfredo; Ramón y Cajal University Hospital, Medical Oncology Department Real, Francisco; Centro Nacional de Investigaciones Oncológicas, Programa de Patología Molecular; Furlong, Laura; Research Program on Biomedical Informatics (GRIB), Hospital del Mar Research Institute (IMIM), Universidad Pompeu Fabra (UPF) Malats, Nuria; Spanish National Cancer Research Center (CNIO), Genetic and Molecular Epidemiology Group; Spanish National Cancer Research Centre (CNIO), Genetic and Molecular Epidemiology Group
Key Words:	Pancreatic cancer risk, Autoimmune diseases, Multimorbidity, Genetic network, Gene-disease associations, Case-control study

SCHOLARONE™
Manuscripts

Short title: Autoimmune diseases and pancreatic cancer risk

Corresponding author: Núria Malats, M.D., M.P.H., Ph. D. Genetic and Molecular

Epidemiology group, Spanish National Cancer Research Center (CNIO), C/Melchor

Fernandez Almagro 3, 28029, Madrid, Spain. Phone: +34-912-246-900 (ext. 3330), Fax:

+34-912-246-911, E-mail: nmalats@cnio.es

Key words: Pancreatic cancer risk; Autoimmune diseases; Multimorbidity; Genetic network; Gene-disease associations; Case-control study.

Abbreviations: Autoimmune disease (AID), Biological Process (BP), Disease Specificity (DSI), Gene-disease association (GAD), Pancreatic cancer (PC), Disease Pleiotropy (DPI).

Article category: Research article

Novelty and impact: Using a bioinformatics approach we show that autoimmune diseases share genetic components with pancreatic cancer and further corroborate this association in a European case-control study population. Some of these results are confirmatory of the mechanisms of pancreatic carcinogenesis while others might point to novel mechanisms. This information could open new venues to explore and increase our understanding of PC risk, potentially impacting the prevention and treatment strategies for this deadly cancer.

Title: Pancreatic cancer and autoimmune diseases: An association sustained by computational and epidemiological case-control approaches

Authors: Paulina Gomez-Rubio, PhD (1,2); Janet Piñero, PhD (3); Esther Molina-Montes, PhD (1,2); Alba Gutiérrez-Sacristán, PhD (3); Mirari Marquez, PhD (1,2); Marta Rava, PhD (1,2); Christoph W. Michalski, MD PhD (4,5); Antoni Farré, MD, PhD (6); Xavier Molero, MD, PhD (2,7,8); Matthias Löhr, MD, PhD (9); José Perea, MD PhD (10); William Greenhalf, PhD (11); Michael O’Rorke, PhD (12); Adonina Tardón, MD, PhD (2,13); Thomas Gress, MD, PhD (14); Victor M. Barberà, PhD (15); Tatjana Crnogorac-Jurcevic, PhD (16); Luís Muñoz-Bellví, MD (17); Enrique Domínguez-Muñoz, MD (18); Joaquim Balsells, MD (2,7,8); Eithne Costello, PhD (11); Jingru Yu (9); Mar Iglesias, MD (19); Lucas Ilzarbe, MD (19); Jörg Kleeff, MD, PhD (4,20); Bo Kong, PhD (4); Josefina Mora, PhD (6); Liam Murray, PhD (12); Damian O’Driscoll, PhD (21); Ignasi Poves, MD (19); Rita T. Lawlor, PhD (22); Weimin Ye, MD, PhD (23); Manuel Hidalgo, MD, PhD (24,25); Aldo Scarpa, MD, PhD (22); Linda Sharp, PhD (21,26); Alfredo Carrato, MD, PhD (2,27); Francisco X. Real* (2,28,29); Laura I. Furlong* (3); Núria Malats (1,2); on behalf of the PanGenEU Study Investigators (30).

* Equal contributions

Authors’ affiliations

- 49 (1) Genetic and Molecular Epidemiology Group, Spanish National Cancer Research Center
50 (CNIO), Madrid, Spain.
- 51 (2) Centro de Investigación Biomédica en Red en Oncología (CIBERONC), Enfermedades
52 Hepáticas y Digestivas (CIBERHD), and Epidemiología y Salud Pública (CIBERESP),
53 Spain.
- 54 (3) Research Program on Biomedical Informatics (GRIB), Hospital del Mar Research
55 Institute (IMIM), Universidad Pompeu Fabra (UPF), Barcelona, Spain.
- 56 (4) Technical University of Munich, Department of Surgery, Munich, Germany.
- 57 (5) University of Heidelberg, Department of Surgery, Heidelberg, Germany.
- 58 (6) Hospital de la Santa Creu i Sant Pau, Department of Gastroenterology, Barcelona, Spain.
- 59 (7) Hospital Universitari Vall d'Hebron, Exocrine Pancreas Research Unit and Vall
60 d'Hebron Research Institute (VHIR), Barcelona, Spain.
- 61 (8) Universitat Autònoma de Barcelona, Campus de la UAB, Barcelona Spain.
- 62 (9) Karolinska Institutet and University Hospital, Gastrocentrum Stockholm, Sweden.
- 63 (10) University Hospital 12 de Octubre, Department of Surgery, Madrid, Spain.
- 64 (11) The Royal Liverpool University Hospital, Department of Molecular and Clinical Cancer
65 Medicine, Liverpool, UK.
- 66 (12) Queen's University Belfast, Centre for Public Health, Belfast, UK.
- 67 (13) Instituto Universitario de Oncología del Principado de Asturias, Department of
68 Medicine, Oviedo, Spain.
- 69 (14) University Hospital of Giessen and Marburg, Department of Gastroenterology, Marburg,
70 Germany.
- 71 (15) Hospital General Universitario de Elche, Laboratorio de Genética Molecular, Elche,
72 Spain.

(16) Barts Cancer Institute, Queen Mary University of London, Centre for Molecular
Oncology, John Vane Science Centre, London, UK.

(17) Hospital Universitario de Salamanca, General and Digestive Surgery Department,
Salamanca, Spain.

(18) Hospital Clínico Universitario de Santiago de Compostela, Department of
Gastroenterology, Santiago de Compostela, Spain.

(19) Hospital del Mar/Parc de Salut Mar, Department of Gastroenterology, Barcelona, Spain.

(20) Department of Visceral, Vascular and Endocrine Surgery, Martin-Luther-University
Halle-Wittenberg, Halle (Saale), Germany.

(21) National Cancer Registry Ireland, Cork, Ireland.

(22) University Hospital Trust of Verona, ARC-Net Centre for Applied Research on Cancer
and Department of Pathology and Diagnostics, Verona, Italy.

(23) Karolinska Institutet and University Hospital, Department of Medical Epidemiology and
Biostatistics, Sweden.

(24) Hospital Madrid-Norte-Sanchinarro and Spanish National Cancer Research Centre
(CNIO), Madrid, Spain.

(25) Rosenberg Clinical Cancer Center, Beth Israel Deaconess Medical Center, Harvard
Medical School, Boston, US.

(26) Institute of Health and Society, Newcastle University, UK.

(27) Hospital Ramón y Cajal, Department of Oncology, Madrid, Spain.

(28) Epithelial Carcinogenesis Group, Spanish National Cancer Research Centre (CNIO),
Madrid, Spain.

(29) Universitat Pompeu Fabra, Departament de Ciències Experimentals i de la Salut,
Barcelona, Spain.

(30) PanGenEU Study Investigators (Additional file 1: Annex S1)

98

99

100

101 **Word count**

102 Abstract: 245 words

103 Main text: 3,426 words

104 Number of figures: 1

105 Number of tables: 4

106 Additional file 1: Annex, 4 Tables, 3 Figures

107

108

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Deciphering the underlying genetic basis behind pancreatic cancer (PC) and its associated multimorbidities will enhance our knowledge towards PC control. The study investigated the common genetic background of PC and different morbidities through a computational approach and further evaluated the less explored association between PC and autoimmune diseases (AIDs) through an epidemiological analysis. Gene-disease associations (GDAs) of 26 morbidities of interest and PC were obtained using the DisGeNET public discovery platform. The association between AIDs and PC pointed by the computational analysis was confirmed through multivariable logistic regression models in the PanGen European case-control study population of 1,705 PC cases and 1,084 controls. Fifteen morbidities shared at least one gene with PC in the DisGeNET database. Based on common genes, several AIDs were genetically associated with PC pointing to a potential link between them. An epidemiologic analysis confirmed that having any of the nine AIDs studied was significantly associated with a reduced risk of PC (Odds Ratio (OR)=0.74, 95% Confidence Interval (CI) 0.58-0.93) which decreased in subjects having ≥ 2 AIDs (OR=0.39, 95%CI 0.21-0.73). In independent analyses, polymyalgia rheumatica and rheumatoid arthritis were significantly associated with low PC risk (OR=0.40, 95%CI 0.19-0.89, and OR=0.73, 95%CI 0.53-1.00, respectively). Several inflammatory-related morbidities shared a common genetic component with PC based on public databases. These molecular links could shed light into the molecular mechanisms underlying PC development and simultaneously generate novel hypotheses. In this study, we report sound findings pointing to an association between AIDs and a reduced risk of PC.

135

136 Introduction

137
138 Pancreatic cancer (PC) is a deadly disease¹ and projected to become the second cause of
139 cancer-related death in Europe by 2030², supporting its consideration as a public health
140 emergency³. Patients with chronic pancreatitis, type 2 diabetes (T2D), or obesity have an
141 increased risk of PC⁴⁻⁶ while those with nasal allergies and asthma have a reduced risk⁷.
142 Multiple morbidities are common in ageing, a main risk factor of PC, but they are seldom
143 studied in a contextual frame, which is important since multiple conditions can share adverse
144 lifestyle and/or genetic susceptibility. In this respect, a higher risk of PC has been reported
145 among subjects having ≥ 3 metabolic syndrome-related disorders or gastric conditions versus
146 those having none of them^{8,9}. Understanding the mechanisms shared between
147 multimorbidities and PC could help improve primary prevention, early-diagnosis, prognosis
148 and/or treatment of PC. Therefore, the combined use of well-annotated epidemiological and
149 clinical datasets with bioinformatics tools becomes ideal to systematically explore the genetic
150 complexity underlying the associations between different multimorbidities and PC.

151 Autoimmune diseases (AIDs) are characterized by an immune dysregulation in which
152 immune cells react against self-antigens resulting in cell and tissue damage. AIDs are
153 classified into organ-specific and systemic depending on whether the autoimmune response is
154 directed against a single or multiple tissues¹⁰. Cancer is thought to result from the
155 accumulation of genetic alterations and the evasion of the immune response against
156 neoantigens; therefore, it is conceivable that AIDs and cancer share genetic mechanisms.
157 Studies have reported positive associations between overall cancer risk and AIDs such as
158 celiac disease, Crohn's disease, rheumatoid arthritis, or systemic lupus erythematosus¹¹⁻¹³.
159 However, contrary associations have been reported for specific types of cancer, e.g. positive
160 associations have been observed for Non-Hodgkin's lymphoma with Crohn's disease,

1
2
3 161 rheumatoid arthritis, and lupus erythematosus, or liver cancer with celiac disease, ulcerative
4
5 162 colitis, or inflammatory bowel disease ^{12,14–17}, while other studies have reported negative
6
7 163 associations between breast cancer with celiac disease, rheumatoid arthritis, and lupus
8
9 164 erythematosus, or colorectal cancer with rheumatoid arthritis^{18–21}. However, to date, limited
10
11 165 and conflicting information exists regarding the association between autoimmunity and the
12
13 166 risk of PC, which could be partly explained by the relatively low prevalence of both
14
15 167 conditions in the population ^{12,17,18,22,23}.

16
17
18 168 In this study, we examine the genetic background shared between 26 candidate
19
20 169 medical conditions and PC to identify underlying common genes using DisGeNET, a
21
22 170 platform that integrates information on gene-disease associations from different public
23
24 171 resources including the literature. Our aim was to generate novel genetic-based hypothesis
25
26 172 regarding multimorbidities associated with PC as well as to obtain insight into their
27
28 173 underlying molecular mechanisms. Finally, we sustain these observations through an
29
30 174 epidemiological approach using the resources of a large international case-control study.

31
32
33 175

34
35 176 **Material and Methods**

36 177
37 178 The bioinformatics analysis was performed with disgenet2r version 3.1.2
38
39 179 (https://bitbucket.org/ibi_group/disgenet2r), an R package that explores the molecular basis
40
41 180 of comorbidities based on DisGeNET²⁴ (version 3.0), a knowledge platform on human
42
43 181 diseases and their association with gene alterations reported in UniProt, ClinVar, and CTD
44
45 182 datasources.

46
47
48 183 *Disease vocabulary.* To interrogate the DisGeNET database, diseases were defined as
49
50 184 Concept Unique Identifiers (CUIs) from the Unified Medical Language Systems
51
52 185 Metathesaurus. The CUIs for PC and 26 candidate morbidities were selected from three
53
54 186 semantic types (disease or syndrome, sign or symptom and neoplastic process) by two

members of the group (PG and NM). A mean of 4 (range 1 to 12) CUIs were used to define each disease (Additional file 1: Table S1). All 26 candidate morbidities were included in the bioinformatics analysis based on their availability of epidemiological information in the PanGenEU study.

DisGeNET query. For all but four conditions, the curated subset of DisGeNET was queried. Polymyalgia rheumatica, mumps, pernicious anaemia, and Addison's disease were not found in the curated data sources and thus were queried from all sources that include data automatically extracted from the literature using text mining approaches. The obtained genes were manually curated by a member of the group (PG).

Disease association on the basis of shared genes. The number of common genes between disease pairs was used to determine disease "genetic" similarities. The Jaccard index (JI) was calculated to estimate the association among diseases accounting for variation in gene findings due differentially studied morbidities, i.e. diseases with more versus less total number of genes identified. The JI is defined as: $|\text{Genesdis1} \cap \text{Genesdis2}| / |\text{Genesdis1} \cup \text{Genesdis2}|$, where Genesdis1 and Genesdis2 are the genes associated with disease 1 and 2, respectively, \cap is the intersection operator, and \cup is the union operator between the two sets of genes. An empirical P value was calculated for each JI using 50,000 bootstrapped samples from a pool of the 7,878 disease genes available in DisGeNET (curated data set). Additionally, the Disease Specificity (DSI) and Pleiotropy (DPI) indexes were calculated (www.disgenet.org/web/DisGeNET/menu/dbinfo#specificity) to characterize all disease genes. Both indexes range from 0 to 1, where DSI = 1 implies high disease specificity (genes associate only with one disease) and DPI = 1 implies high disease pleiotropy (the gene is associated with several diseases and these belong to different classes).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Gene Ontology analysis. The Gene Ontology terms, restricted to the Biological Process (BP) branch, were identified through an enrichment analysis performed with the common genes between disease pairs with GO.db and GOSTats packages in R.

Pathway analysis. An enrichment analysis was performed using the ReactomePA package in R to identify pathways shared between morbidities based on common genes between disease pairs.

The epidemiological analysis was performed using resources from the PanGenEU case-control study conducted in Spain, United Kingdom, Germany, Ireland, Sweden, and Italy, between 2009-2014 (Additional file 1: Annex S1).

Study population. Potentially eligible pancreatic ductal adenocarcinoma patients, men and women ≥ 18 years of age, were approached for participation. Subjects with diagnosis not confirmed by the physician were excluded from the study. Eligible controls did not have previous history of PC and their primary diagnosis for hospital admission was not associated with known risk factors of PC (Supplementary Table S1 and S2) when recruited in a hospital-based setting; controls from Ireland and Sweden were population-based. Subjects without information in the entire medical section of the questionnaire were excluded (N=268, 8.8%) leaving 1,705 PC cases and 1,084 controls for analyses. Age, sex, and smoking distributions were similar between included and excluded subjects ($P > 0.05$). An overall response rate of 86.3% was observed for cases and 77.8% for controls. IRB ethical approval and written informed consent was obtained by all participating centers and study participants, respectively.

Study variables. Demographics, lifestyle, and medical history were gathered through in-person interviews applying standardized questionnaires. Subjects reported “yes/no” to “Has your doctor ever told you that you had any of the following illnesses, health problems or procedures?” for 26 candidate morbidities including 9 AIDs further categorized as

systemic (rheumatoid arthritis, lupus erythematosus, scleroderma, and polymyalgia rheumatica), hematologic (pernicious anaemia), gastrointestinal (Crohn's disease, ulcerative colitis, and celiac disease), and endocrine (Addison's disease). Regular consumption of anti-inflammatory/pain killer medication (aspirin, paracetamol, NSAIDs and corticosteroids) was defined as subjects reporting ever taking one of these treatments at least once a week, on average, for 3 months or more.

Statistical analysis. Imputation of missing values (3.7% in cases and 2.9% in controls) was performed with random forest (missForest R package). Missing values were assumed to be missing at random. Variables used for imputation (% missings) included case-control status, country, age (2.3%), sex (0.2%), smoking (pack/years, 9.4%), alcohol status (1.5%), medication (6.1- 7.9%), and morbidities and time since diagnosis (0.8%-21.7%). Imputation was performed with no maximum number of iterations and 100 trees. An imputation test introducing the same proportions of missing values to a complete-case dataset resulted in a concordance mean between imputed and real data >90% and out of bag errors < 0.35 (0= good imputation performance, 1= bad imputation performance). Multivariable logistic regression models were used to test the association between PC risk and AIDs, individually and by defined groups. Adjustment variables were selected based on the 10% change in estimate, the likelihood ratio tests and the Akaike Information Criterion (AIC). Potential confounders considered were smoking, alcohol, T2D, obesity, family history of PC, years of education, and treatment. Multicollinearity between variables was discarded based on variance-inflation factor threshold <2. Interactions with age, sex, smoking, alcohol and treatments were explored by including the interaction terms in the models.

Results

1
2
3 261 The gene-disease association analysis based on publicly available information showed that all
4
5 262 diseases except hypothyroidism, were connected through shared genes with at least one other
6
7 263 morbidity (Additional file 1: Figure S1). The total number of genes associated with the 26
8
9 264 non-cancer medical conditions averaged 41.3 ranging from 1 in heartburn to 185 in
10
11 265 hypertension (Table 1). The strongest associations based on common genes were found
12
13 266 between mumps and polymyalgia, Crohn’s disease and ulcerative colitis, asthma and nasal
14
15 267 allergies, and diabetes and hypertension (average JI= 0.124, range, 0.10–0.14) (Additional
16
17 268 file 1: Table S3). The average number of biological processes and pathways shared between
18
19 269 morbidities was 147.2 (range 9-386) and 29.7 (range 4-96).

20
21
22 270 DisGeNET curated subset showed 73 genes associated with PC; half of these had
23
24 271 disease specificity >0.75 and disease pleiotropy <0.30 (Additional file 1: Table S4). Twenty-
25
26 272 two out of the 73 genes (range 0-10) were also associated with other morbidities (Tables 1
27
28 273 and 2), with *ABO*, *SPINK1*, *PDX1*, *TFPI2*, and *STK11* showing the highest disease specificity
29
30 274 and the lowest disease pleiotropy (DSI > 0.73 and DPI < 0.41).

31
32
33 275 Fifteen morbidities were associated with PC through at least one gene, including
34
35 276 peptic ulcer (JI=0.055), hypertension (JI=0.04), and ulcerative colitis (JI=0.04) that showed
36
37 277 the most robust genetic relationship with PC (Table 1 and Figure 1). PC and hypertension had
38
39 278 the highest number of shared biological processes (N=69); 125 unique biological processes
40
41 279 were shared between PC and seven morbidities (Table 3). The biological process that was
42
43 280 shared between PC and more morbidities was ‘negative regulation of fat cell differentiation’.
44
45 281 Twenty-one unique pathways were shared between PC and at least one morbidity with peptic
46
47 282 ulcer sharing the highest number of pathways (N=16) (Table 3). ‘TNF signalling’ and
48
49 283 ‘TNFR1-mediated proapoptotic signalling’ pathways were shared between PC, T2D, peptic
50
51 284 ulcer, and rheumatoid arthritis.
52
53
54
55
56
57
58
59
60

285 Giving the unexpected genetic link observed between AIDs and PC in DisGeNET and
286 the existing scarce literature regarding this relationship ^{4,9,25}, we decided to focus on these
287 conditions in the PanGenEU epidemiological study and assessed the association with PC risk
288 of nine AIDs with enough available data in the study population. Overall, 16.2% of controls
289 reported having any AID in comparison to 13% reported by PC cases (Table 4).
290 Multivariable logistic regression models showed that having any AID was associated with a
291 significant reduced risk of PC (OR=0.74, 95%CI 0.58-0.93). Furthermore, the number of
292 AIDs was significantly associated with a reduced PC risk trend (P trend = 0.002). Having any
293 systemic or organ-specific AID was significantly associated with lower PC risk (OR=0.74,
294 95%CI: 0.55-0.99, and OR=0.71, 95%CI 0.52-0.97, respectively). Among organ-specific
295 diseases, having any one or more gastrointestinal AID was borderline associated with a low
296 risk of PC (OR= 0.51, 95%CI 0.26-1.00). Analysis of individual AIDs showed significant
297 and borderline significant associations with polymyalgia rheumatica and rheumatoid arthritis,
298 respectively (OR=0.4, 95%CI: 0.18-0.89, and OR=0.73, 95%CI: 0.53-1.00). The association
299 with any AID and ≥ 2 AIDs was maintained after inflammatory/pain killer treatment
300 adjustment. No significant interactions were observed.

301

302 Discussion

303 We used a systems medicine approach to unravel the shared genetic background of PC and its
304 associated co-morbidities. We show that out of the 26 morbidities of interest 15, including
305 five AIDs, share a genetic background with PC according to the available data in DisGeNET,
306 a fact that points to the involvement of immune dysregulation processes in PC pathogenesis.
307 We confirmed the AID-PC association first observed in DisGeNET with the results from a
308 large European case-control study population.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

We observed that some morbidities previously associated with PC risk in the literature^{9,26} were also sustained at a genetic level sharing 22 genes with PC according to DisGeNET. Many of the genes highlighted in these explorations are relevant in pancreatic carcinogenesis (e.g. *KRAS*, *ABO*, *BRCA2*, *STK11*), while others are important players in inflammatory process (e.g. *PTGS2*, *TGFB1*, *CXCL8*)^{27,28}. Likewise, many of the biological processes and pathways shared between these morbidities and PC are essential in inflammation and carcinogenesis.

Disentangling the underlying mechanisms behind multimorbidities could improve disease prevention and patient management. Growing evidence suggests that integrative approaches help to better grasp disease complexity. New bioinformatics tools such as DisGeNET that generate multimorbidity networks based on gene-disease associations available from public datasets have become ideal for this purpose. A main limitation of this approach is that we can only consider gene-disease associations that have been published, curated and registered in databases. In our study, the few or lack of genetic associations between some morbidities may result from either a real absence of a genetic association or due to the incomplete knowledge about the genetic basis of human diseases. Therefore, we cannot rule out that some associations could have been missed due to the still limited spectrum of the information available. In this respect, we relied on the JI to overcome problems due to differences in how well studied some diseases are over others. Conversely, we cannot exclude the possibility that other morbidities not included in our bioinformatics analysis could be also genetically associated with PC. Assessing the common genetic background of other autoimmune morbidities and PC might help us to gather an even broader view of this malignancy and further increase our understanding of PC.

Applying a systems approach to explore PC-associated multimorbidities has the potential to generate new hypotheses. Among the 15 morbidities found genetically associated

with PC, rheumatoid arthritis, lupus erythematosus, ulcerative colitis, Crohn's disease, and scleroderma have been vaguely studied as risk factors of PC (Additional file 1: Figure S2). Previous work suggested that AIDs could predispose individuals to cancer. Additionally, paraneoplastic autoimmune syndromes have been described²⁹. Current studies point to an association between certain AIDs and an increased risk of different types of cancers such as small intestine and oesophageal cancers with celiac disease, or haematological neoplasms with rheumatoid arthritis, systemic lupus erythematosus, and scleroderma^{11,17,19,30}. Yet, some of these AIDs have been associated with a decreased risk of other cancers such as breast and colorectal^{18,19,31}, suggesting that the association may not be the same for all cancer types. For PC, the information is limited and conflicting.

The association with pernicious anaemia seems to be the only one that has been purposely analysed in the context of PC, but no significant association was reported³². Other studies exploring the association between specific AIDs and cancer overall mainly relying on registry data and reporting on PC, appear inconclusive^{22,33–35}. The retrospective nature of most of these studies limits the adjustment for potential confounders. Moreover, to our knowledge, no study explored the association between PC and scleroderma. Our study design allowed us to account for confounding and interaction of a broad set of factors.

We report a significant negative association between suffering one or more of the nine AIDs under study and PC risk. We observed that the estimate further decreased in subjects reporting ≥ 2 AIDs, with a significant negative trend. Independent analysis of each AID showed a significant association between polymyalgia rheumatica and rheumatoid arthritis with lower PC risk. However, these analyses were limited by sample size. Other studies have reported a lack of significant association between lupus or Addison's disease and PC^{13,17,20,22,23,31,36}. Contradictory results have been published regarding pernicious anaemia, Crohn's disease, and ulcerative colitis with some studies reporting no significant

1
2
3 359 association^{16,22,32,33,35} while others showing a significant increased PC risk^{12,15,34}. For celiac
4
5 360 disease, studies have reported both significantly reduced and increased risk of PC^{18,22}, but a
6
7 361 recent meta-analysis showed no significant association¹¹. Furthermore, most studies have
8
9 362 described a lack of association between rheumatoid arthritis and PC risk^{22,37-39}. However, a
10
11 363 study performed with the Scottish Cancer Registry reported a significant negative association
12
13 364 between rheumatoid arthritis and PC among women but not among men²¹; in our study, no
14
15 365 interaction was observed with gender. Other studies evaluating ulcerative colitis, Crohn's
16
17 366 disease, rheumatoid arthritis, polymyalgia and giant cell arthritis, and pernicious anaemia by
18
19 367 time since diagnosis reported significantly increased PC risk mostly restricted to subjects
20
21 368 reporting <1 year difference between AIDs and cancer diagnoses; when >1 year between
22
23 369 diagnoses is reported, loss of significance and smaller PC risk estimates are commonly
24
25 370 observed^{12,15,34,40,41}. ~~The latter results in combination with the results obtained in this study~~
26
27 371 ~~suggest a negative association between PC risk and AIDs.~~ Comparison with the existing
28
29 372 literature is challenging since, to our knowledge, this is the first study incorporating
30
31 373 confounders and combining different AIDs into single variables. Although the
32
33 374 epidemiological literature is inconclusive regarding their link, atopy and autoimmunity are
34
35 375 known hypersensitive reactions of the immune system⁴². Accordingly, our enrichment
36
37 376 analyses showed that atopic and autoimmune diseases share several immune-related
38
39 377 biological processes and pathways (Additional file 1: Figure S3). The importance of the
40
41 378 immune environment in which tumours develop is further strengthened by the emerging
42
43 379 evidence of the role of immune checkpoint inhibition as a potent therapeutic strategy in a
44
45 380 wide variety of tumours. However, only a fraction of patients respond to such therapies and
46
47 381 there is an urgent need to identify predictors of response⁴³. These results, and our previous
48
49 382 findings of a reduced PC risk among subjects with atopic conditions⁷, strongly suggest that
50
51
52
53
54
55
56
57
58
59
60

immune recognition of neoantigens, its quantity and quality could contribute to modulate the immune response to emerging preneoplastic clones⁴⁴.

Conversely, while autoimmune pancreatitis is an AID suggested to develop into chronic pancreatitis, its link with PC risk has not been established^{45,46}. Moreover, it is reasonable to expect a different type of association in AIDs that result in localized damage to the pancreas. Moreover, one could hypothesize that the decreased risk of certain cancers and AIDs could be the consequence of confounding by particular treatments. However, we lacked information about treatment for AIDs. *In vivo* studies suggest that anti-TNF therapy could inhibit pancreatic tumour growth and metastasis^{47,48}, though its link with carcinogenesis is not completely elucidated^{49,50}. Combination of gemcitabine with TNF inhibitors at doses approved for AID treatment does not seem to significantly improve gemcitabine treatment in PC patients⁵¹. We show that further adjusting for anti-inflammatory/pain relief treatment results in the loss of significance for some associations. While these results could suggest a potential confounding effect, additionally supporting common pathways between PC and AID, we cannot completely rule out chance findings giving the small effect of the treatment adjustment on the estimates and the sustained significance of the main variables.

The interpretation and generalization of these results must be done carefully. While the combination of AIDs into categories helped us to overcome the problem of statistical power, our study size is limited for the inquiry of specific disorders and stratified analysis. Additionally, considering that our information is self-reported, some degree of disease misclassification might have occurred resulting in a lower rate of false positive than false negative reports, probably those with less severe conditions, which would attenuate the risk estimates. It remains necessary to replicate these associations; in this context register-based information and electronic Medical Records could be an ideal setting to gather a longitudinal view of the association between AIDs and PC.

1
2
3 408
4
5 409
6
7 410
8
9 411
10
11 412
12
13 413
14
15 414
16
17 415
18
19 416
20
21 417
22
23
24 418
25

Conclusion

To our knowledge this is the first study exploring in depth the association between several autoimmune diseases and PC risk. This report highlights the importance of multimorbidities in PC risk. Many of these results are confirmatory of the mechanistic notions about pancreatic carcinogenesis. Importantly, common genetic association between PC and AIDs was identified through a bioinformatics approach, which was further characterized in an epidemiological setting as a negative association between AIDs and PC risk, opening new venues to explore and increase our understanding of PC risk potentially impacting the prevention and treatment strategies for this deadly cancer.

26 419
27
28 420
29
30 421
31
32 422
33
34 423
35
36 424
37
38 425

Acknowledgements. We acknowledge the coordinators, field and administrative workers, technicians, and study participants of the European Study into Digestive Illnesses and Genetics (PanGenEU) study and the COST Action BMI1204 ‘An integrated European platform for pancreatic cancer research: from basic science to clinical and public health interventions for a rare disease’.

39 425
40
41 426
42
43 427
44
45 428
46
47 429
48
49 430
50
51 431
52
53 432
54
55 432
56
57
58
59
60

Funding. The work was partially supported by Fondo de Investigaciones Sanitarias (FIS), Instituto de Salud Carlos III-FEDER, Spain (#PI11/01542, #PI0902102, #PI12/01635, #PI12/00815, #PI13/00082, CP10/00524, PI15/01573); Red Temática de Investigación Cooperativa en Cáncer, Spain (#RD12/0036/0034, #RD12/0036/0050, #RD12/0036/0073); EU H2020 Programme 2014-2020 (634143 MedBioinformatics, and 676559 Elixir-Excelerate); World Cancer Research (WCR #15-0391); Acción Especial de Genómica, Spain (#GEN2001-4748-c05-03); EU-6FP Integrated Project (#018771-MOLDIAG-PACA); EU-FP7-HEALTH (#259737-CANCERALIA, #256974-EPC-TM-Net, #602783-Cam-Pac);

Cancer Focus Northern Ireland and Department for Employment and Learning; ALF (#SLL20130022), Sweden; Associazione Italiana Ricerca Cancro [grant number: 12182]; FP7 European Community Grant Cam-Pac [no: 602783]; Italian Ministry of Health [FIMPCUP_J33G13000210001].

Author's contributions. Paulina Gómez-Rubio and Núria Malats had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Study concept and design: Paulina Gomez-Rubio, Francisco X Real, Laura I Furlong, Núria Malats. Acquisition, analysis, or interpretation of data: All authors. Drafting the manuscript: Paulina Gomez-Rubio, Janet Piñero, Laura I Furlong, Núria Malats. Critical revision of the manuscript for important intellectual content: All authors. Statistical and bioinformatics analyses: Paulina Gomez-Rubio, Janet Piñero. Obtained funding: Christoph Michalski, Matthias Löhr, William Greenhalf, Thomas Gress, Tatjana Crnogorac-Jurcevic, Liam Murray, Rita Lawlor, Aldo Scarpa, Linda Sharp, Francisco X Real, Laura I Furlong, Núria Malats. Study supervision: Núria Malats, Francisco X Real, Laura I Furlong.

Conflict of interests. The authors declare none.

References

1. De Angelis R, Sant M, Coleman MP, Francisci S, Baili P, Pierannunzio D, Trama A, Visser O, Brenner H, Ardanaz E, Bielska-Lasota M, Engholm G, et al. Cancer survival in Europe 1999-2007 by country and age: results of EURO CARE-5-a population-based study. *Lancet Oncol* 2013;2045:1–12.

1
2
3 458 2. Rahib L, Smith BD, Aizenberg R, Rosenzweig AB, Fleshman JM, Matrisian LM.
4
5 459 Projecting Cancer Incidence and Deaths to 2030: The Unexpected Burden of Thyroid,
6
7 460 Liver, and Pancreas Cancers in the United States. *Cancer Res* 2014;74:2913–21.
8
9
10 461 3. L  hr JM. Pancreatic cancer should be treated as a medical emergency. *BMJ*
11
12 462 2014;349:g5261.
13
14 463 4. Raimondi S, Lowenfels AB, Morselli-Labate AM, Maisonneuve P, Pezzilli R.
15
16 464 Pancreatic cancer in chronic pancreatitis; aetiology, incidence, and early detection.
17
18 465 *Best Pract Res Clin Gastroenterol* 2010;24:349–58.
19
20
21 466 5. Bosetti C, Rosato V, Li D, Silverman D, Petersen GM, Bracci PM, Neale RE, Muscat
22
23 467 J, Anderson K, Gallinger S, Olson SH, Miller AB, et al. Diabetes, antidiabetic
24
25 468 medications, and pancreatic cancer risk: an analysis from the International Pancreatic
26
27 469 Cancer Case-Control Consortium. *Ann Oncol* 2014;25:2065–72.
28
29
30 470 6. Aune D, Greenwood DC, Chan DSM, Vieira R, Vieira a R, Navarro Rosenblatt D a,
31
32 471 Cade JE, Burley VJ, Norat T. Body mass index, abdominal fatness and pancreatic
33
34 472 cancer risk: a systematic review and non-linear dose-response meta-analysis of
35
36 473 prospective studies. *Ann Oncol* 2012;23:843–52.
37
38
39 474 7. Gomez-Rubio P, Zock J-P, Rava M, Marquez M, Sharp L, Hidalgo M, Carrato A,
40
41 475 Ilzarbe L, Michalski C, Molero X, Farr   A, Perea J, et al. Reduced risk of pancreatic
42
43 476 cancer associated with asthma and nasal allergies. *Gut* 2015;gutjnl-2015-310442.
44
45
46 477 8. Rosato V, Tavani A, Bosetti C, Pelucchi C, Talamini R, Polesel J, Serraino D, Negri E,
47
48 478 La Vecchia C. Metabolic syndrome and pancreatic cancer risk: A case-control study in
49
50 479 Italy and meta-analysis. *Metabolism* 2011;60:1372–8.
51
52
53 480 9. Gomez-Rubio P, Rosato V, Marquez M, Bosetti C, Molina-Montes E, Rava M, Pi  nero
54
55 481 J, Michalski CW, Farr   A, Molero X, L  hr M, Ilzarbe L, et al. A systems approach

- identifies time-dependent associations of multimorbidities with pancreatic cancer risk.
Ann Oncol 2017;[Epub ahead of print].
10. Fairweather D. Autoimmune Disease: Mechanisms. *Encycl Life Sci* 2007;1–6.
11. Han Y, Chen W, Li P, Ye J. Association Between Coeliac Disease and Risk of Any Malignancy and Gastrointestinal Malignancy: A Meta-Analysis. *Medicine (Baltimore)* 2015;94:e1612.
12. Hemminki K, Li X, Sundquist J, Sundquist K. Cancer risks in Crohn disease patients. *Ann Oncol* 2009;20:574–80.
13. Mellemkjær L, Andersen V, Linet MS, Gridley G, Hoover R, Olsen JH. Non-Hodgkin's lymphoma and other cancers among a cohort of patients with systemic lupus erythematosus. *Arthritis Rheum* 1997;40:761–8.
14. Askling J, Linet M, Gridley G, Halstensen TS, Ekström K, Ekbom A. Cancer incidence in a population-based cohort of individuals hospitalized with celiac disease or dermatitis herpetiformis. *Gastroenterology [Internet]* 2002;123:1428–35. Available from: <http://www.sciencedirect.com/science/article/pii/S0016508502002755>
15. Hemminki K, Li X, Sundquist J, Sundquist K. Cancer risks in ulcerative colitis patients. *Int J Cancer* 2008;123:1417–21.
16. Pedersen N, Duricova D, Elkjaer M, Gamborg M, Munkholm P, Jess T. Risk of extra-intestinal cancer in inflammatory bowel disease: meta-analysis of population-based cohort studies. *Am J Gastroenterol* 2010;105:1480–7.
17. Cao L, Tong H, Xu G, Liu P, Meng H, Wang J, Zhao X, Tang Y, Jin J. Systemic lupus erythematosus and malignancy risk: A meta-analysis. *PLoS One* 2015;10:1–21.
18. Ilus T, Kaukinen K, Virta LJ, Pukkala E, Collin P. Incidence of malignancies in

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

505 diagnosed celiac patients: a population-based estimate. *Am J Gastroenterol*
506 2014;109:1471–7.

507 19. Simon T, Thompson A, Gandhi K, Hochberg M, Suissa S. A meta-analysis of the
508 incidence of malignancy in adult patients with rheumatoid arthritis. *Arthritis Res Ther*
509 2015;17:212.

510 20. Parikh-Patel A, Allen M, Cress R, White RH. Cancer risk in a cohort of patients with
511 systemic lupus erythematosus (SLE) in California. *Cancer Causes Control*
512 2008;19:887–94.

513 21. Thomas E, Brewster DH, Black RJ, Macfarlane GJ. Risk of malignancy among
514 patients with rheumatic conditions. *IntJCancer* 2000;88:497–502.

515 22. Landgren AM, Landgren O, Gridley G, Graça M, Linet MS, Morton LM. Autoimmune
516 disease and subsequent risk of developing alimentary tract cancers among 4.5 million
517 U.S. male Veterans. 2011;117:1163–71.

518 23. Dreyer L, Faurschou M, Mogensen M, Jacobsen S. High incidence of potentially
519 virus-induced malignancies in systemic lupus erythematosus: A long-term followup
520 study in a Danish cohort. *Arthritis Rheum* 2011;63:3032–7.

521 24. Piñero J, Queralt-Rosinach N, Bravo Á, Deu-Pons J, Bauer-Mehren A, Baron M, Sanz
522 F, Furlong LI. DisGeNET: A discovery platform for the dynamical exploration of
523 human diseases and their genes. *Database* 2015;2015:1–17.

524 25. Maisonneuve AP, Amar S, Lowenfels AB. Periodontal Disease , Edentulism and
525 Pancreatic Cancer : A Meta Analysis. *Ann Oncol* 2017;28:985–95.

526 26. Maisonneuve P, Lowenfels a. B. Risk factors for pancreatic cancer: a summary review
527 of meta-analytical studies. *Int J Epidemiol* 2014;1–13.

- 528 27. Amundadottir LT. Pancreatic cancer genetics. *Int J Biol Sci* 2016;12:314–25.
- 529 28. Hausmann S, Kong B, Michalski C, Erkan M, Friess H. The Role of Inflammation in
530 Pancreatic Cancer. *Adv Exp Med Biol* 2014;816:129–51.
- 531 29. Rahat MA, Shakya J. Parallel Aspects of the Microenvironment in Cancer and
532 Autoimmune Disease. *Mediators Inflamm* 2016;2016:4375120.
- 533 30. Bonifazi M, Tramacere I, Pomponio G, Gabrielli B, Avvedimento E V, La Vecchia C,
534 Negri E, Gabrielli A. Systemic sclerosis (scleroderma) and cancer risk: systematic
535 review and meta-analysis of observational studies. *Rheumatology (Oxford)*
536 2013;52:143–54.
- 537 31. Bernatsky S, Ramsey-Goldman R, Labrecque J, Joseph L, Boivin JF, Petri M, Zoma
538 A, Manzi S, Urowitz MB, Gladman D, Fortin PR, Ginzler E, et al. Cancer risk in
539 systemic lupus: An updated international multi-centre cohort study. *J Autoimmun*
540 2013;42:130–5.
- 541 32. Shah P, Rhim A, Haynes K, Hwang W, Yang Y. Diagnosis of pernicious anemia and
542 the risk of pancreatic cancer. *Pancreas* 2014;19:161–9.
- 543 33. Brinton LA, Gridley G, Hrubec Z, Hoover R, Fraumeni JF. Cancer risk following
544 pernicious anaemia. *Br J Cancer* 1989;59:810–3.
- 545 34. Hsing AW, Hansson L-E, McLaughlin JK, Nyren O, Blot WJ, Ekblom A, Fraumeni JF.
546 Pernicious anemia and subsequent cancer. A population-based cohort study. *Cancer*
547 1993;71:745–50.
- 548 35. Møller M, Gridley G, Møller H, Hsing AW, Linet MS, Brinton LA, Olsen JH.
549 Pernicious anaemia and cancer risk in Denmark. *Br J Cancer* 1996;73:998–1000.
- 550 36. Abu-Shakra M, Gladman DD, Urowitz MB. Malignancy in systemic lupus

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

551 erythematosus. *Arthritis Rheum* 1996;39:1050–4.

552 37. Gridley G, McLaughlin JK, Ekblom A, Klareskog L, Adami HO, Hacker DG, Hoover
553 R, Fraumeni JF. Incidence of cancer among patients with rheumatoid arthritis. *J Natl*
554 *Cancer Inst* 1993;85:307–11.

555 38. Askling J, Fored CM, Brandt L, Baecklund E, Bertilsson L, Feltelius N, Cöster L,
556 Geborek P, Jacobsson LT, Lindblad S, Lysholm J, Rantapää-Dahlqvist S, et al. Risks
557 of solid cancers in patients with rheumatoid arthritis and after treatment with tumour
558 necrosis factor antagonists. *Ann Rheum Dis* 2005;64:1421–6.

559 39. Yu K-H, Kuo C-F, Huang LH, Huang W-K, See L-C. Cancer Risk in Patients With
560 Inflammatory Systemic Autoimmune Rheumatic Diseases: A Nationwide Population-
561 Based Dynamic Cohort Study in Taiwan. *Medicine (Baltimore)* 2016;95:e3540.

562 40. Hemminki K, Li X, Sundquist K, Sundquist J. Cancer risk in hospitalized rheumatoid
563 arthritis patients. *Rheumatology* 2008;47:698–701.

564 41. Ji J, Liu X, Sundquist K, Sundquist J, Hemminki K. Cancer risk in patients
565 hospitalized with polymyalgia rheumatica and giant cell arteritis: A follow-up study in
566 Sweden. *Rheumatology* 2010;49:1158–63.

567 42. Rabin RL, Levinson AI. The nexus between atopic disease and autoimmunity: A
568 review of the epidemiological and mechanistic literature. *Clin Exp Immunol*
569 2008;153:19–30.

570 43. Turnis ME, Andrews LP, Vignali DAA. Inhibitory receptors as targets for cancer
571 immunotherapy. *Eur J Immunol* 2015;45:1892–905.

572 44. Balachandran VP, Luksza M, Zhao JN, Makarov V, Moral JA, Remark R, Herbst B,
573 Askan G, Bhanot U, Senbabaoglu Y, Wells DK, Cary CIO, et al. Identification of
574 unique neoantigen qualities in long-term survivors of pancreatic cancer. *Nature*

- 2017;551:S12–6.
45. Maruyama M, Watanabe T, Kanai K, Oguchi T, Asano J, Ito T, Ozaki Y, Muraki T, Hamano H, Arakura N, Kawa S. Autoimmune pancreatitis can develop into chronic pancreatitis. *Orphanet J Rare Dis* 2014;9:77.
46. Hart PA, Law RJ, Dierkhising RA, Smyrk TC, Takahashi N, Chari ST. Risk of cancer in autoimmune pancreatitis: A case-control study and review of the literature. *Pancreas* 2014;43.
47. Egberts JH, Cloosters V, Noack A, Schniewind B, Thon L, Klose S, Kettler B, von Forstner C, Kneitz C, Tepel J, Adam D, Wajant H, et al. Anti-tumor necrosis factor therapy inhibits pancreatic tumor growth and metastasis. *Cancer Res* 2008;68:1443–50.
48. Tiwari S, Egberts JH, Kornienko O, Köhler L, Trauzold A, Glüer CC, Kalthoff H. Assessment of anti-inflammatory tumor treatment efficacy by longitudinal monitoring employing sonographic micro morphology in a preclinical mouse model. *BMC Med Imaging* 2011;11:15.
49. Nyboe Andersen N, Pasternak B, Basit S, Andersson M, Svanstrom H, Caspersen S, Munkholm P, Hviid A, Jess T. Association between tumor necrosis factor-alpha antagonists and risk of cancer in patients with inflammatory bowel disease. *JAMA* 2014;311:2406–13.
50. Shelton E, Laharie D, Scott FI, Mamtani R, Lewis JD, Colombel J, Ananthakrishnan AN. Cancer Recurrence Following Immune-Suppressive Therapies. *Gastroenterology* 2016;151:97–109.e4.
51. Wu C, Fernandez S, Criswell T, Chidiac T, Guttridge D, Villalona-Calero M, Bekaii-Saab T. Disrupting cytokine signaling in pancreatic cancer: a phase I/II study of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

599 etanercept in combination with gemtacinibine in patients with advanced disease.
600 *Pancreas* 2013;42:813–8.
601
602

For Peer Review

603

Table 1. Number of concept unique identifiers, total genes, common genes with pancreatic cancer and respective Jaccard indexes for diseases under study.

Disease1	Number of CUIs	Genes _{dis1}	Common Genes	JI	P val ^a
Peptic ulcer	3	23	5	0.0549	< 10 ⁻⁵
Hypertension	9	185	10	0.0403	< 10 ⁻⁵
Ulcerative colitis	1	57	5	0.04	< 10 ⁻⁵
Acid regurgitation	2	5	2	0.0263	< 10 ⁻⁵
Asthma	6	83	5	0.0331	0.0001
Obesity	10	156	7	0.0315	0.0001
Type 2 diabetes	3	124	6	0.0314	0.0002
<i>H.pylori</i>	2	5	1	0.013	0.0009
Rheumatoid arthritis	1	123	5	0.0262	0.001
Chronic pancreatitis	1	6	1	0.0128	0.0013
Periodontitis	4	9	1	0.0123	0.0032
Hyperthyroidism	4	10	1	0.0122	0.0037
Scleroderma	12	11	1	0.012	0.0048
Crohn's disease	3	41	2	0.0179	0.0065
Lupus erythematosus	6	68	2	0.0144	0.0250
Nasal allergies	3	17	0	0	-
Skin allergies	9	43	0	0	-
Hypercholesterolemia	12	40	0	0	-
Heartburn	1	1	0	0	-
Hypothyroidism	2	3	0	0	-
Mumps	1	9	0	0	-
Pernicious anaemia	1	2	0	0	-
Gallstones	3	6	0	0	-
Celiac disease	1	28	0	0	-
Polymyalgia	3	7	0	0	-
Addison disease	1	11	0	0	-

^aEmpirical p value calculated after performing 50,000 bootstraps

CUI: Concept unique identifier, JI= Jaccard index

Note: pancreatic cancer CUIs = 3, Genes = 73.

604

605

606

607

1
2
3 608
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36 609
37
38 610
39
40 611
41
42
43 612
44
45 613
46
47
48 614
49
50
51 615
52
53 616
54
55
56 617
57
58
59
60

Table 2. Common genes between pancreatic cancer and other diseases in DisGeNET.

Gene	N _{dis}	Diseases
TNF	9	Asthma, obesity, type 2 diabetes, hypertension, peptic ulcer, rheumatoid arthritis, scleroderma, Crohn's disease, ulcerative colitis
MMP9	6	Asthma, obesity, peptic ulcer, periodontitis, lupus erythematosus, ulcerative colitis
PTGS2	6	Obesity, hypertension, peptic ulcer, acid regurgitation, rheumatoid arthritis, lupus erythematosus
SOD2	5	Obesity, type 2 diabetes, hypertension, hyperthyroidism, rheumatoid arthritis,
PPARG	4	Obesity, type 2 diabetes, hypertension, Crohn's disease
TGFB1	4	Asthma, type 2 diabetes, hypertension, peptic ulcer
AHR	2	Hypertension, rheumatoid arthritis
CDH1	2	Ulcerative colitis, <i>H. pylori</i> infection
CXCL8	2	Acid regurgitation, Ulcerative colitis
PTEN	2	Asthma, hypertension
ABO	1	Peptic ulcer
BCL2L1	1	Type 2 diabetes
CNR1	1	Obesity
DPYD	1	Obesity
HIF1A	1	Hypertension
PDX1	1	Type 2 diabetes
PLAU	1	Asthma
SPINK1	1	Chronic pancreatitis
STAT3	1	Ulcerative colitis
STK11	1	Hypertension
TFPI2	1	Rheumatoid arthritis
TP53	1	Hypertension

618

Table 3. Number of unique biological processes and pathways shared between each disease and any other diseases and number of processes shared between the diseases and pancreatic cancer.

Disease	N unique processes annotated	N processes shared with PC	N unique pathways annotated	N pathways shared with PC
Hypertension	386	69	96	0
Type 2 diabetes	340	18	54	10
Asthma	278	14	46	0
Rheumatoid arthritis	277	13	36	7
Crohn's disease	266	0	36	0
Obesity	251	31	37	0
Ulcerative colitis	195	18	30	0
Peptic ulcer	177	31	37	16
Lupus erythematosus	169	0	32	0
Skin allergies	139	0	26	0
Pancreatic cancer	125	-	21	-
Hyperthyroidism	50	0	12	0
Hypercholesterolemia	36	0	16	0
Celiac disease	30	0	12	0
Mumps	24	0	4	0
Addison disease	17	0	14	0
Polymyalgia	17	0	16	0
Scleroderma	10	0	6	0
Nasal allergies	9	0	0	0

PC: pancreatic cancer

Note: Only diseases for which biological processes and/or pathways were found through enrichment analyses are presented.

619

620

621

622

623

624

Table 4. Odds ratios for the association between autoimmune diseases and pancreatic ductal adenocarcinoma.

	Cases		Controls		OR ^a	95%CI	OR ^b	95%CI	OR ^c	95%CI
	N=1705	%	N=1084	%						
Autoimmune diseases										
No	1483	86.9	908	83.8	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.
Any	222	13.0	176	16.2	0.78	[0.62;0.98]	0.74	[0.58;0.93]	0.78	[0.61;0.99]
Number of AID										
No	1483	86.9	908	83.8	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.
1	201	11.8	146	13.5	0.85	[0.66;1.08]	0.81	[0.63;1.04]	0.85	[0.66;1.09]
≥2	21	1.2	30	2.8	0.45	[0.25;0.83]	0.39	[0.21;0.73]	0.44	[0.24;0.81]
<i>P trend^d</i>					0.009		0.002		0.01	
Type of AID										
None	1483	86.9	908	83.8	Ref.	Ref.				
Systemic only	120	7.0	86	7.9	0.85	[0.62;1.16]	0.78	[0.58;1.08]	0.83	[0.60;1.15]
Organ-specific only	88	5.2	73	6.7	0.77	[0.55;1.08]	0.76	[0.54;1.07]	0.78	[0.55;1.11]
Hematologic	65	3.8	53	4.9	0.82	[0.56;1.21]	0.80	[0.54;1.18]	0.82	[0.55;1.21]
Gastrointestinal	20	1.2	15	1.4	0.58	[0.25;1.32]	0.60	[0.26;1.37]	0.64	[0.28;1.46]
Endocrine	2	0.1	1	0.1	1.84	[0.16;2.11]	2.65	[0.23;3.07]	2.91	[0.25;3.34]
Mixed organ-specific	1	0.1	4	0.4	0.28	[0.03;2.64]	0.29	[0.03;2.74]	0.29	[0.03;2.85]
Both systemic and localized	14	0.8	17	1.6	0.50	[0.24;1.07]	0.42	[0.19;0.90]	0.48	[0.22;1.05]
Systemic AID										
No	1571	92.1	981	90.5	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.
Any	134	7.9	103	9.5	0.81	[0.60;1.08]	0.74	[0.55;0.99]	0.79	[0.59;1.07]
Organ-specific AID										
No	1603	94	994	91.7	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.
Any	102	6	90	8.3	0.73	[0.53;0.99]	0.71	[0.52;0.97]	0.74	[0.54;1.02]
Gastrointestinal AID										
No	1679	98.5	1059	97.7	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.
Any	26	1.5	25	2.3	0.51	[0.26;0.99]	0.51	[0.26;1.00]	0.55	[0.28;1.09]
Lupus										
No	1696	99.5	1081	99.7	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.
Yes	9	0.5	3	0.3	1.91	[0.48;7.53]	2.08	[0.51;8.44]	2.04	[0.49;8.39]
Scleroderma										
No	1697	99.5	1080	99.6	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.
Yes	8	0.5	4	0.4	0.91	[0.23;3.54]	0.81	[0.20;3.20]	0.83	[0.21;3.25]
Polymyalgia										
No	1692	99.2	1065	98.3	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.

Yes	13	0.8	19	1.8	0.42	[0.20;0.92]	0.40	[0.18;0.89]	0.46	[0.21;1.04]
Pernicious anemia										
No	1630	95.6	1014	93.5	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.
Yes	75	4.4	70	6.5	0.75	[0.53;1.06]	0.72	[0.51;1.02]	0.75	[0.53;1.07]
Crohn's disease										
No	1701	99.8	1079	99.5	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.
Yes	4	0.2	5	0.5	0.26	[0.05;1.46]	0.32	[0.05;1.94]	0.36	[0.58;2.24]
Celiac disease										
No	1698	99.6	1078	99.5	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.
Yes	7	0.4	6	0.6	0.54	[0.15;1.95]	0.63	[0.18;2.17]	0.62	[0.18;2.19]
Addison's disease										
No	1698	99.6	1078	99.5	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.
Yes	7	0.4	6	0.6	0.89	[0.30;2.68]	0.91	[0.29;2.81]	0.96	[0.31;2.99]
Rheumatoid arthritis										
No	1597	93.7	1000	92.3	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.
Yes	108	6.3	84	7.8	0.81	[0.59;1.11]	0.73	[0.53;1.00]	0.78	[0.56;1.08]
Ulcerative colitis										
No	1690	99.1	1068	98.5	Ref.	Ref.	Ref.	Ref.	Ref.	Ref.
Yes	15	0.9	16	1.5	0.55	[0.24;1.30]	0.49	[0.21;1.16]	0.54	[0.23;1.30]

^a Adjusted for age (continuous), sex and country

^b Model adjusted for age (continuous), sex, country, smoking (pack/years, tertiles based on the population distribution and alcohol status (never, former, current)

^c Model adjusted for age (continuous), sex, country, smoking (pack/years, tertiles based on the population distribution, alcohol status (never, former, current), and treatment (no treatment, aspirin only, NSAIDs only, paracetamol only, corticosteroids only, and more than one treatment type)

^d Trend was calculated using number of AID as a continuous variable (range from 0 to 4)

Figure 1. Gene network of medical conditions associated with PC through common genes. A) Network of diseases that share genes with pancreatic cancer and all corresponding connections; B) Network of diseases that share genes with pancreatic cancer, only connections with pancreatic cancer shown. Edge width represents the Jaccard index for each disease pair; Jaccard indexes were multiplied by 100 in order to allow better visualization. Node size represents the number of genes obtained through DisGeNET for each medical condition.




Figure 1. Gene network of medical conditions associated with PC through common genes. A) Network of diseases that share genes with pancreatic cancer and all corresponding connections; B) Network of diseases that share genes with pancreatic cancer, only connections with pancreatic cancer shown. Edge width represents the Jaccard index for each disease pair; Jaccard indexes were multiplied by 100 in order to allow better visualization. Node size represents the number of genes obtained through DisGeNET for each medical condition.

37x48mm (300 x 300 DPI)